

B4



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C07K 14/81, C12N 15/52, 15/10, C12Q 1/68</b>		A1	(11) International Publication Number: <b>WO 00/11034</b>
			(43) International Publication Date: 2 March 2000 (02.03.00)
(21) International Application Number: PCT/EP99/06054 (22) International Filing Date: 18 August 1999 (18.08.99) (30) Priority Data: 98115957.7                      25 August 1998 (25.08.98)                      EP (71)(72) Applicant and Inventor: KLEESIEK, Knut [DE/DE]; Goethestrasse 14, D-32545 Bad Oeynhausen (DE). (72) Inventors; and (75) Inventors/Applicants (for US only): BRINKMANN, Thomas [DE/DE]; Estelistrasse 9, D-33739 Bielefeld (DE). PRO- HASKA, Wolfgang [DE/DE]; Brekerfeld 1, D-32547 Bad Oeynhausen (DE). GOETTING, Christian [DE/DE]; Bahn- hofstrasse 84, D-32584 Loehne (DE). SCHMIDT, Michael [DE/DE]; Schlosshofstrasse 213, D-33615 Bielefeld (DE). (74) Agent: BENZ, Juergen; Heumann, Benz, Spessarttring 63, D-64287 Darmstadt (DE).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>	
(54) Title: A MUTANT OF TISSUE FACTOR PATHWAY INHIBITOR, DNA SEQUENCE AND USE FOR DETECTING THROM- BOTIC DISORDERS  (57) Abstract  The invention relates to a novel mutant of tissue factor pathway inhibitor (TFPI) protein and its corresponding DNA sequence. The mutant can be found in humans who show or may show an increased risk of thrombotic diseases. By screening samples of human blood for said DNA or fragments of it, it is possible to predict a disposition of thrombotic disorders by which prophylactic application or measures can be initiated.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## **A Mutant of Tissue Factor Pathway Inhibitor, DNA Sequence and Use for Detecting Thrombotic Disorders**

5 The invention relates to a novel mutant of tissue factor pathway inhibitor (TFPI) protein and its corresponding DNA sequence. The mutant can be found in humans who show or may show an increased risk of thrombotic diseases. The DNA sequence according to this invention differs from the known TFPI coding DNA by a single nucleotide polymorphism which results in a change of one amino  
10 acid position within the known TFPI protein. By screening samples of human blood for said DNA or fragments of it, it is possible to predict a disposition of thrombotic disorders by which prophylactic application or measures can be initiated.

15 Tissue factor pathway inhibitor (TFPI) is an important regulator in the extrinsic blood coagulation pathway. Although the regulatory biochemical role of TFPI is evident, the clinical significance of this proteinase inhibitor remains to be elucidated. The definition of a clinical TFPI deficiency seems to be more complex than that of other coagulation inhibitors because the activity and concentration of  
20 circulating TFPI can not be considered a true measure of *in vivo* levels. Its determination in plasma samples by immunological methods or functional assays has been shown to be inadequate in the detection of a clinical deficiency.

TFPI is a single chain glycoprotein present in plasma in trace amounts. It was  
25 previously known as extrinsic pathway inhibitor, or lipoprotein associated coagulation inhibitor. TFPI belongs to the class of Kunitz-type proteinase inhibitors, and the mature protein contains an acidic amino-terminal end followed by three Kunitz-type inhibitory domains and a basic carboxy-terminal end. The cDNA coding for TFPI was cloned and characterized by Wun et al. (J. Biol.  
30 Chem. (1988), 263). The mature molecule consists of 276 amino acid residues, including 18 cysteins (see Seq. Id. No. 1, 2), all involved in disulfide bonds, and contains three potential N-linked glycosylation sites. The molecular weight of the polypeptide backbone is about 32 kDa; the protein present in plasma runs, however, on SDS-PAGE with an apparent molecular weight of about 42 kDa,

presumably due to glycosylation.

The multivalent protease inhibitor is an important regulator of the extrinsic pathway of blood coagulation through its ability to interact with the blood coagulation factor VIIa / tissue factor complex and the activated factor X via its

- 5 Kunitz-type domains  $\delta 1$  and  $\delta 2$  (see: Girard, T.J. et al. , *Nature* **338**, 518-520 (1989); Broze, G.J.Jr. et al.; *Blood* **71**, 335 (1984); Rapaport, S.I. & Rao, L.V.M., *Thrombosis and Haemostasis* **74**, 7-17 (1995); Broze, G.J.Jr. , *Haemostaseologie* **17**, 73-77 (1997) ). There is also evidence that infusion of recombinant TFPI may protect against disseminated intravascular coagulation induced by TF or E. coli to
- 10 protect against venous thrombosis and to prevent rethrombosis after successful thrombolysis in arterial thrombosis. The intravascular distribution of TFPI is complex. The mature human tissue factor pathway inhibitor protein is mainly synthesized by the vascular endothelium (Bajaj, M.S. et al. *Proc. Natl. Acad. Sci. USA* **88**, 8869, (1990)). It has also been detected in at least four intravascular
- 15 pools: bound to the endothelial cell surface, associated with lipoproteins, carrier-free within the plasma, and sequestered in platelets (Sandset, P.M. & Abildgaard, U., *Haemostasis* **21**, 219 (1991) ).

A review of the regulation and role of TFPI within the extrinsic pathway system is given by Petersen et al. (*Thrombosis Research*, **79**, 1-47 (1995)).

20

- TFPI plays such an important role in the inhibition of the extrinsic pathway that TFPI deficiencies due to mutations in the TFPI gene should enhance the activity of the prothrombinase complex. This increases the thrombin generation and consequently the risk of venous thrombosis. Such a diminished inhibition of
- 25 thrombin generation is already well known in inherited coagulation inhibitor defects that predispose to thrombosis including deficiency of antithrombin III, protein C and protein S (Dahlbäck, B. *Blood* **85**, 607-614 (1995) ). The most prevalent inherited abnormality which is known to lead to venous thrombosis is the resistance to activated protein C caused by a single point mutation in the
- 30 factor V gene (Bertina, R.M. et al., *Nature* **369**, 64-67 (1994)).

Thus it was an object of the present invention to screen genomic DNA samples of human normal blood donors and thrombotic patients for alterations in the TFPI gene to assess the influence of a modified TFPI in venous thromboembolic

diseases.

Surprisingly, a single nucleotide substitution (C→T) in exon 7 of the TFPI gene could be detected, leading to a proline to leucine exchange at amino acid position 151 of the mature protein (see Seq. Id. No. 3, 4). Investigating the statistical  
5   significance of this modification of TFPI it was found that said modification can be linked to a relative risk to general thrombotic disorders. However, it should be pointed out that said modification seems also to occur in human individuals showing evidently - by comparing their family health history - no increased risk to thrombotic events. Nevertheless, the finding of modified TFPI protein / DNA in  
10   blood samples of a human individual may provide an important hint for a possible disposition of said individual for thrombotic diseases. Therefore, prophylactic measures can be taken in order to prevent said diseases.

According to this invention the term "thrombotic disorders" used above and below  
15   includes all known diseases or malfunctions which can be related directly or indirectly to a permanent or temporary abnormal or pathological blood coagulation, e.g. venous thrombosis.

The modified TFPI protein or related DNA can be used as diagnostic marker to  
20   detect such a possible risk to thrombotic disorders in a patient.

Thus, it is an object of this invention to provide a DNA sequence coding for a mutant of tissue factor pathway inhibitor (TFPI) protein, wherein a proline residue at position 151 of the mature peptide (calculated from the N-terminus) is replaced  
25   by a leucine residue.

It is a further object of the invention to provide a DNA sequence comprising a DNA sequence (a) coding for a signal peptide followed by a DNA sequence (b) coding for a mutant of TFPI protein, wherein a cytosine at position 536, calculated  
30   from the start codon of the sequence (a) is replaced by a tyrosine to form a CTG codon instead of a CCG codon within the coding region of sequence (b).

Furthermore, it is an object of the invention to provide a diagnostic method to detect a disposition for thromboembolic diseases in humans by screening

genomic DNA samples of human blood for a DNA sequence defined above or in the claims.

Especially, the invention relates to a diagnostic method using polymerase chain reaction (PCR) and restriction analysis by extracting total DNA from human blood samples amplifying exon 7 of the TFPI gene ( from position 536 to 628) and its previous flanking intronic region by means of suitable primers, treating the PCR products with a restriction enzyme having the recognition site ACTGG or CAGTG, and detecting the fragment length after restriction analysis or isolating and detecting the DNA defined above and in the claims.

10

Suitable primers according to this inventions are primers or DNA fragments which may hybridize with the corresponding regions of exon 7 of the TFPI gene, intronic flanking regions included. Therefore, as preferred embodiments, it is an object of this invention to provide a diagnostic method defined above and in the claims using the following primers (Seq. Id. No. 6 , 8):

5' - TCTATTTTAATTGGCTGTAT - 3' and

5' - GCATGATAATAGTTTCCTGG - 3'.

The single nucleotide polymorphism modification (C → T, CCG → CTG) in the TFPI gene at position 536 creates a new cleavage site within this region of the gene, which is not present in the original gene. This is very advantageous and can be preferably used for short evidence of such modified DNA in blood samples and, as consequence, of the above-mentioned disposition for thrombotic disorders. The recognition site created by said nucleotide exchange is ACTGG or CAGTG (Seq. ID. No. 5, 7). Therefore, all restriction enzymes which can recognize this cleavage site are suitable to carry out the diagnostic method of the invention.

Known and suitable restriction enzymes are, for example, Bse1I, BseNI, BsrI, BsrSI, BscH1, Bst11I, BsoHI, Tsp1I, and TspRI. The preferred restriction enzyme of the invention is BseNI.

Finally, it is an object of the present invention to provide the novel mutant of tissue factor pathway inhibitor (TFPI) protein wherein a proline residue at position 151 of the mature peptide (calculated from the N-terminus, without signal peptide

sequence) is replaced by a leucine residue.

According to this invention it was proved the thesis that genetic variations in the TFPI gene contribute to the occurrence of hitherto unexplained cases of thrombophilia in about half of the afflicted patients. In a first screening experiment  
5 50 unrelated individuals with a thrombotic history were investigated, who were selected to determine the genetic basis of their thrombosis (27 of them were shown to be carriers of the factor V Leiden mutation). While scanning all coding exons of the TFPI gene and the adjacent 5' and 3' intronic regions by PCR-SSCP analysis, an abnormal pattern suggesting the presence of a genetic variation, was  
10 observed in a PCR fragment from exon 7 (Fig. 1a). DNA sequencing of the fragment showing the abnormal SSCP banding pattern revealed a single heterozygous C to T mutation at nucleotide position 1 of exon 7, changing the codon CCG<sup>151</sup> to CTG<sup>151</sup> resulting in a Pro<sup>151</sup> to Leu<sup>151</sup> exchange in the amino acid of the mature protein (Fig. 1b, c; Fig. 2) (see also: *van der Logt et al.*,  
15 *Biochemistry* **30**, 1571-1577 (1991); *Girard, T.J. et al.*; *J. Biol. Chem.* **266**, 5036-5041 (1991)).

The 536C→T transition is associated with the creation of a new recognition site for the restriction enzyme *BseNI*, providing a rapid means of screening further  
20 individuals for this mutation by PCR and restriction fragment length polymorphism analysis (PCR-RFLP). The primers designed for the amplification of exon 7 and the 5' and 3' flanking intronic regions were used to generate a 170 bp PCR fragment. If the nucleotide C is present at position 536, the 170 bp DNA fragment is not digestible with *BseNI*. However, if the nucleotide T is present at this  
25 position a 27 bp and a 143 bp restriction fragment is generated. Using this method a second set of patients was screened with venous thrombosis (n=324) of whom 30.2% carried the factor V Leiden mutation. Another individual heterozygous for the TFPI mutation was detected. Homozygous carriers of this mutation were not found.

30

In order to estimate the prevalence of the 536C→T exchange in the general population, 2480 randomly chosen unrelated blood donors (age 18 - 60 years) were investigated by PCR-RFLP analysis.

To detect a greater number of subjects with the trait, which would allow a more precise estimation of the relative risk of those individuals to develop venous thrombosis, family members of the heterozygote blood donors and patients were investigated. In total 13 heterozygote individuals within 6 different families were  
5 found. Two of them suffered from deep vein thrombosis.

All subjects carrying the TFPI mutation were investigated for the presence of other genetic defects of clotting proteins (factor V Leiden, prothrombin 20210G→A mutation, deficiency of protein C, protein S and antithrombin III) in  
10 order to exclude the contribution of these disorders to the thromboembolic findings. The two individuals who showed the TFPI trait and suffered from venous thrombosis had none of the additionally examined genetic disorders. Although the TFPI mutation was observed together with either the factor V Leiden mutation or the prothrombin mutation (20210G→A) in 4 members of one family, none of them  
15 had a history of thromboembolic diseases.

It is generally accepted that direct evidence for an important regulatory role of an inhibitory clotting protein requires the detection of low levels of the circulating inhibitor associated with venous thrombosis. Therefore, the TFPI activity was  
20 measured by a functional assay and the protein concentration by an immunological assay in plasma of all individuals showing the TFPI mutation. Compared with the control group (blood donors without TFPI mutation) no statistically significant differences were detected. However, existing small differences might not have been recognized, due to the low number of cases  
25 investigated so far.

Furthermore, for an interpretation of plasma TFPI levels it must be taken into consideration that compared with other coagulation inhibitors such as antithrombin III and protein C which circulate mainly as free molecules, circulating  
30 TFPI may not reflect the true measure of the total *in vivo* TFPI quantity. Therefore, determination of TFPI activity or protein concentration in plasma samples has been shown to be an inadequate approach to detect clinical TFPI deficiencies. The intravascular distribution of TFPI is more complex. A major pool of TFPI (about 50-80% of the total intravascular pool) is normally bound to the



endothelium, but may be released into the circulation following injection of heparin. It is also a known fact that more than 80% of the circulating TFPI is in a complex form with lipoprotein, preferentially with low density lipoprotein. However, whether this fraction of lipoprotein-associated TFPI remains active as a clotting inhibitor is still unknown. The plasma TFPI level correlates with the lipoprotein concentration, and it has been shown that patients with an inherited abetalipoproteinemia have decreased TFPI plasma concentrations but do not suffer from venous thrombosis. Therefore, the questions arises whether a deficiency of total intravascular TFPI can be detected by any of the tests used so far. (Sandset, P.M. & Bendz, B. *Thrombosis and Haemostasis* 78, 467-470 (1997)). A genetic abnormality of TFPI which promotes the development of venous thrombosis may affect different functions of the protein, such as secretion by endothelial cells, binding to the endothelial membrane, proteolysis in the vascular space, and association with lipoproteins.

So far, 13 individuals with the TFPI mutation were identified. Two of them had a history of venous thrombosis. Although the number of subjects is rather small to evaluate the risk of thrombophilia, the prevalence of venous thromboembolism in this group was compared with the prevalence in 466 blood donors who were thoroughly questioned about a possible history of venous thrombosis. 10 cases were found in this group. Statistical analysis showed a probability of 94.5% for the hypothesis that the presence of the TFPI trait is linked to thrombophilia ( $p=0.055$ ; odds ratio: 5.9; 95% confidence interval: 1.0-36.5). Thus, this question remains open and can only be answered when more subjects with the TFPI mutation are available for statistical evaluation.

#### **Description of the figures:**

**Fig. 1:** PCR-SSCP and sequence change for the proline to leucine substitution in a TFPI variant.

- a) Single strand conformation polymorphism in exon 7 of the TFPI gene in a thrombotic patient (lane C) and nonthrombotic controls (lanes A, B, D). The arrow in lane C indicates an additional band only observed in heterozygous variants.
- b) Sequencing profile of the sense strand of the PCR products obtained from total DNA from the heterozygous person identified by SSCP. The C→T transition in

the nucleotide sequence, indicated by two peaks at the same position in the heterozygous probe is marked by an arrow.

- c) The C→T transition at nucleotide position 1 of exon 7 results in a CCG → CTG change in the sense strand and leads to a proline to leucine substitution at position 151 of the TFPI protein.

**Fig. 2:** Proposed secondary structure of the tissue factor pathway inhibitor and the position of the amino acid exchange. The mature protein consists of 276 amino acids, forming three tandem *Kunitz*-type proteinase inhibitory domains, two connecting chains, an acidic N-terminus with negatively charged amino acids (-) and a basic carboxy-terminal end, containing a cluster of positively charged amino acids (+). *Kunitz*-domain  $\delta 1$  has been identified as binding site for the factor VIIa / tissue factor complex, while domain  $\delta 2$  binds activated factor Xa (Wesselschmidt, R. et al., *Blood* **79**, 2004-2010 (1992)). Whether *Kunitz*-type domain  $\delta 3$  binds and inhibits a specific protease is unknown. The basic carboxy-terminal end is the predicted binding site to glycosaminoglycans at the endothelial cell surface and contributes to heparin binding (Enjyoji, K.-I. et al., *Biochemistry* **34**, 5725-5735 (1995)). Truncated forms of TFPI are tightly bound to low density lipoproteins and lack the distal portion of the full-length molecule including *Kunitz*-type domain  $\delta 3$  (Broze, G.J.Jr. et al., *Blood Coag. Fibrinol.* **5**, 551-559 (1994)). Asn<sup>117</sup> and Asn<sup>167</sup> are N-glycosylated, Ser<sup>174</sup> and Thr<sup>175</sup> are O-glycosylated (\*) (Nakahara, T.M. et al. *Biochemistry* **35**, 6450-6459 (1996)). The Pro<sup>151</sup> to Leu change is located near the *Kunitz*-type inhibitor domain  $\delta 2$  within the second connecting chain (magnification).

25

The following examples describe the invention in more detail without restricting it.

#### Example 1:

*Polymerase chain reaction (PCR).* 374 unrelated patients with a definite history of deep vein thrombosis and 2480 blood donors as a control group were investigated in this study. Total DNA of all subjects was extracted from whole blood using the QIAamp Blood Kit (QIAGEN, Hilden, Germany). TFPI-exon 7 and its 5' and 3' flanking intronic regions were specifically amplified from genomic DNA by PCR (Saiki, R.K. et al., *Science* **239**, 487-491 (1988)). The primers for

the amplification reaction (TFPI Ex7F 5'-TCTATTTTAATTGGCTGTAT-3', TFPI Ex7R 5'-GCATGATAATAGTTTCCTGG-3') were derived from the genomic sequence of the TFPI gene (*van der Logt et al., Biochemistry* **30**, 1571-1577 (1991)). The standard PCR included 0.1 - 1 µg of genomic DNA, 300 nM of each primer, 200 µM of each deoxynucleotide triphosphate, GeneAmp 10x PCR Buffer and 2.5 units of AmpliTaq DNA Polymerase (Perkin-Elmer Corporation, Foster City, CA) in a final volume of 50 µl. After mixing, 1 drop of mineral oil was added to each tube to prevent evaporation. Thermal cycling conditions included initial denaturation at 94°C for 3 minutes, followed by 40 cycles of denaturing at 94°C for 30 sec, annealing at 48°C for 45 sec and extending at 72°C for 45 sec. The PCR products were electrophoresed on a neutral 0.8% agarose gel and stained with ethidium bromide for inspection.

#### Example 2:

Single Strand Conformation Polymorphism (SSCP). Sequence variations within the amplified DNA were detected by single strand conformation polymorphism following the PCR. For SSCP analysis the amplified DNA fragments containing the coding region of exon 7 were purified using the QIAquick PCR Purification Kit (QIAGEN, Hilden, Germany). Following this procedure, the fragments were diluted 1 : 6 in distilled water, heated for 10 minutes at 98°C, and then quenched on ice to achieve almost complete denaturation. Electrophoresis was performed using the PhastSystem electrophoresis unit of Amersham Pharmacia Biotech (Uppsala, Sweden). Strand separation was obtained with a 12.5% Homogeneous PhastGel at a temperature of 12°C. Running conditions were: (i) Pre-run: 400 V, 5.0 mA, 1.0 W, 12°C, 60 Vh; (ii) Sample application: 25 V, 5.0 mA, 1.0 W, 12°C; 2 Vh; (iii) Main-run: 200 V, 5.0 mA, 1.0 W; 12°C, 220 Vh. Gels were silver-stained in the coloration unit of the device, following the method described by Bassam *et al.* (Bassam, B.J. *et al., Analyt. Biochem.* **196**, 80-83 (1991)).

#### Example 3:

DNA-Sequence analysis of SSCP variants. The purified PCR fragments showing differences in the SSCP banding patterns were used as template for forward and reverse cycle sequencing reactions with the primers described above. The fragments were sequenced using the Applied Biosystems

Incorporated (ABI) protocol for TAQ cycle sequencing with dye terminators and an automated ABI PRISM 377 DNA Sequencer (Applied Biosystems, Weiterstadt, Germany).

5 **Example 4:**

*PCR - Restriction Fragment Length Polymorphism (RFLP) analysis for the determination of the TFPI 536C → T mutation.*

PCR was carried out as described above. The PCR product was controlled on a 0.8% agarose gel. 6 µl of the PCR product were then incubated for 1.5 h with 5U of *BseNI* (MBI  
10 Fermentas, St. Leon-Rot, Germany) at 65°C in a final volume of 20 µl without further purification. Following this, the samples were loaded onto a 2.5% agarose gel, ethidium bromide stained, and analyzed under UV-light.

**Example 5:**

15 *PCR-RFLP analysis for the determination of the Factor V 1691 G→A mutation.*

The PCR-RFLP analysis of the factor V Leiden mutation was carried out as previously described by Beauchamp *et al.*, using the primers Fv3 and Fv6 (Beauchamp, N.J. *et al.*, *Brit. J. Haematol.* **88**, 219-222 (1994)).

20 **Example 6:**

*PCR-mediated site-directed mutagenesis for the determination of the 20210 G→A transition in the prothrombin gene.*

The G to A transition at position 20210 in the prothrombin gene was determined after amplification with primer PTHF 5'-CGCCTGAAGAAGTGGATACAGA-3' and PTHR 5'-

25 ATAGCACTGGGAGCATTGAA GC-3'. The latter was designed with a C to A substitution at position 20214 to create a restriction site for *HindIII* (MBI Fermentas, St. Leon-Rot, Germany) when the G to A transition at position 20210 is present in the prothrombin gene (Poort, R.S. *et al.*, *Blood* **88**, 3968-3703 (1996)). Restriction analysis and gel electrophoresis were carried out as  
30 described above under reaction conditions recommended by *HindIII*.

**Example 7:**

*Measurement of TFPI concentration and activity in plasma samples.* Total and full-length forms of TFPI, complexes with tissue factor (TF) and factor VIIa as

well as binary complexes with factor Xa and quaternary complexes with TF, factor VIIa and factor Xa were quantified in plasma samples obtained from individuals heterozygous for the TFPI mutation, unaffected members of their families and blood donors, using the IMUBIND Total and Truncated TFPI ELISA Kit (American Diagnostica Inc., Greenwich, CT) according to the manufacturer's instructions. The same plasma samples were used to determine the activity of predominantly free TFPI with the ACTICHROME TFPI Activity Assay from American Diagnostica Inc. (Greenwich, CT).

**Example 8:**

*Determination of protein C and antithrombin III concentration in plasma samples.*

The protein C and AT III concentration in plasma samples was measured using the DADE BEHRING (Liederbach, Germany) Protein C and the Antithrombin III Chromogenic Assays according to the manufacturer's instructions.

15

**Example 9:**

*Determination of protein S activity in plasma samples.* The activity of the protein C cofactor was determined with the Protein S Clotting Test of Boehringer Mannheim (Mannheim, Germany). The clotting times were measured with a ball coagulometer (Amelung, Lemgo, Germany).

20

**Example 10:**

*Determination of high and low density lipoprotein.* To exclude plasma samples from patients with abnormal lipoprotein concentration due to association of TFPI to low density lipoprotein we determined the concentration of high and low density lipoprotein by a commercially available turbidimetric assay (Boehringer Mannheim, Mannheim, Germany).

25

**Example 11:**

*Statistical analysis.* The statistical analysis was performed using Chisquare test and linear logistical regression analysis for sex and age dependent calculations with the Statistical Analysis System (SAS) program, version 6.12 and Student's t-test. For statistical analysis the patients with thrombotic history were matched to the control group according to age and sex.

30

**Patent Claims**

- 5 1. A DNA sequence coding for a mutant of tissue factor pathway inhibitor (TFPI) protein wherein a proline residue at position 151 of the mature peptide (calculated from the N-terminus, without signal peptide) is replaced by a leucine residue.
- 10 2. A DNA sequence comprising a DNA sequence (a) coding for a signal peptide followed by a DNA sequence (b) coding for a mutant of TFPI protein, wherein a cytosine at position 536, calculated from the start codon of the sequence (a) is replaced by a tyrosine to form a CTG codon instead of a CCG codon within the coding region of sequence (b).
- 15 3. A DNA sequence comprising the DNA sequence of Seq. Id. No. 3.
4. Use of a DNA sequence of any of the claims 1 to 3 for the *in vitro* diagnosis of thrombotic disorders.
- 20 5. Use according to claim 4 for the *in vitro* diagnosis of a disposition for venous thromboembolic diseases in humans.
6. A diagnostic method to detect a disposition for thromboembolic diseases in humans by screening genomic DNA samples of human blood for a DNA
- 25 sequence according to any of claims 1 to 3.
7. A diagnostic method of claim 6 using polymerase chain reaction (PCR) and restriction analysis by extracting total DNA from human blood samples
- 30 amplifying exon 7 of the TFPI gene and its previous flanking intronic region by means of suitable primers, treating the PCR products with a restriction enzyme having the recognition site ACTGG or CAGTG, and detecting the fragment length after restriction analysis or isolating and detecting the DNA sequence of any of claims 1 to 3.

8. A diagnostic method of claim 7, wherein the following primers are used:

(a) 5' - TCTATTTTAATTGGCTGTAT - 3' and

(b) 5' - GCATGATAATAGTTTCCTGG - 3'.


5

9. A diagnostic method of any of claims 6 to 8, wherein the restriction enzyme is BseNI.


10. A mutant of tissue factor pathway inhibitor (TFPI) protein wherein a proline  
10 residue at position 151 of the mature peptide (calculated from the N-terminus, without signal peptide) is replaced by a leucine residue.

1/2


A




B



C

Fig. 1



**FIG. 2**

## SEQUENCE LISTING



&lt;170&gt; PatentIn Ver. 2.1

5 <210> 1  
 <211> 912  
 <212> DNA  
 <213> Human

10 <220>  
 <221> CDS  
 <222> (1)..(912)  
 <223> TFPI protein + signal peptide

15 <220>  
 <221> sig\_peptide  
 <222> (1)..(84)  
 <223> lipoprotein-associated coagulation inhibitor

20 <220>  
 <221> mat\_peptide  
 <222> (85)..(912)  
 <223> mature TFPI

25 <400> 1  
 atg att tac aca atg aag aaa gta cat gca ctt tgg gct tct gta tgc 48  
 Met Ile Tyr Thr Met Lys Lys Val His Ala Leu Trp Ala Ser Val Cys  
 -25 -20 -15

30 ctg ctg ctt aat ctt gcc cct gcc cct ctt aat gct gat tct gag gaa 96  
 Leu Leu Leu Asn Leu Ala Pro Ala Pro Leu Asn Ala Asp Ser Glu Glu  
 -10 -5 -1 1

gat gaa gaa cac aca att atc aca gat acg gat ttg cca cca ctg aaa 144  
 35 Asp Glu Glu His Thr Ile Ile Thr Asp Thr Asp Leu Pro Pro Leu Lys  
 5 10 15 20

ctt atg cat tca ttt tgt gca ttc aag gcg gat gat ggc cca tgt aaa 192  
 40 Leu Met His Ser Phe Cys Ala Phe Lys Ala Asp Asp Gly Pro Cys Lys  
 25 30 35

gca atc atg aaa aga ttt ttc ttc aat att ttc act cga cag tgc gaa 240  
 Ala Ile Met Lys Arg Phe Phe Phe Asn Ile Phe Thr Arg Gln Cys Glu  
 40 45 50

45 gaa ttt ata tat ggg gga tgt gaa gaa aat cag aat cga ttt gaa agt 288  
 Glu Phe Ile Tyr Gly Gly Cys Glu Glu Asn Gln Asn Arg Phe Glu Ser  
 55 60 65

50 ctg gaa gag tgc aaa aaa atg tgt aca aga gat aat gca aac agg att 336  
 Leu Glu Glu Cys Lys Lys Met Cys Thr Arg Asp Asn Ala Asn Arg Ile  
 70 75 80

ata aag aca aca ttg caa caa gaa aag cca gat ttc tgc ttt ttg gaa 384  
 55 Ile Lys Thr Thr Leu Gln Gln Glu Lys Pro Asp Phe Cys Phe Leu Glu  
 85 90 95 100

gaa gat cct gga ata tgt cga ggt tat att acc agg tat ttt tat aac 432  
 60 Glu Asp Pro Gly Ile Cys Arg Gly Tyr Ile Thr Arg Tyr Phe Tyr Asn  
 105 110 115

aat cag aca aaa cag tgt gaa cgt ttc aag tat ggt gga tgc ctg ggc 480  
 Asn Gln Thr Lys Gln Cys Glu Arg Phe Lys Tyr Gly Gly Cys Leu Gly  
 120 125 130

5 aat atg aac aat ttt gag aca ctg gaa gaa tgc aag aac att tgt gaa 528  
 Asn Met Asn Asn Phe Glu Thr Leu Glu Glu Cys Lys Asn Ile Cys Glu  
 135 140 145

10 gat ggt cgg aat ggt ttc cag gtg gat aat tat gga acc cag ctc aat 576  
 Asp Gly Pro Asn Gly Phe Gln Val Asp Asn Tyr Gly Thr Gln Leu Asn  
 150 155 160

15 gct gtg aat aac tcc ctg act ccg caa tca acc aag gtt ccc agc ctt 624  
 Ala Val Asn Asn Ser Leu Thr Pro Gln Ser Thr Lys Val Pro Ser Leu  
 165 170 175 180

20 ttt gaa ttt cac ggt ccc tca tgg tgt ctc act cca gca gac aga gga 672  
 Phe Glu Phe His Gly Pro Ser Trp Cys Leu Thr Pro Ala Asp Arg Gly  
 185 190 195

25 ttg tgt cgt gcc aat gag aac aga ttc tac tac aat tca gtc att ggg 720  
 Leu Cys Arg Ala Asn Glu Asn Arg Phe Tyr Tyr Asn Ser Val Ile Gly  
 200 205 210

30 aaa tgc cgc cca ttt aag tac agt gga tgt ggg gga aat gaa aac aat 768  
 Lys Cys Arg Pro Phe Lys Tyr Ser Gly Cys Gly Gly Asn Glu Asn Asn  
 215 220 225

35 ttt act tcc aaa caa gaa tgt ctg agg gca tgt aaa aaa ggt ttc atc 816  
 Phe Thr Ser Lys Gln Glu Cys Leu Arg Ala Cys Lys Lys Gly Phe Ile  
 230 235 240

40 caa aga ata tca aaa gga ggc cta att aaa acc aaa aga aaa aga aag 864  
 Gln Arg Ile Ser Lys Gly Gly Leu Ile Lys Thr Lys Arg Lys Arg Lys  
 245 250 255 260

45 aag cag aga gtg aaa ata gca tat gaa gaa att ttt gtt aaa aat atg 912  
 Lys Gln Arg Val Lys Ile Ala Tyr Glu Glu Ile Phe Val Lys Asn Met  
 265 270 275

<210> 2  
 <211> 304  
 <212> PRT  
 45 <213> Human

<400> 2  
 Met Ile Tyr Thr Met Lys Lys Val His Ala Leu Trp Ala Ser Val Cys  
 -25 -20 -15

50 Leu Leu Leu Asn Leu Ala Pro Ala Pro Leu Asn Ala Asp Ser Glu Glu  
 -10 -5 -1 1

55 Asp Glu Glu His Thr Ile Ile Thr Asp Thr Asp Leu Pro Pro Leu Lys 20  
 5 10 15

Leu Met His Ser Phe Cys Ala Phe Lys Ala Asp Asp Gly Pro Cys Lys  
 25 30 35

60 Ala Ile Met Lys Arg Phe Phe Phe Asn Ile Phe Thr Arg Gln Cys Glu  
 40 45 50

Glu Phe Ile Tyr Gly Gly Cys Glu Glu Asn Gln Asn Arg Phe Glu Ser  
                   55                                  60                                  65  
 5 Leu Glu Glu Cys Lys Lys Met Cys Thr Arg Asp Asn Ala Asn Arg Ile  
           70                                  75                                  80  
 Ile Lys Thr Thr Leu Gln Gln Glu Lys Pro Asp Phe Cys Phe Leu Glu  
       85                                  90                                  95                                  100  
 10 Glu Asp Pro Gly Ile Cys Arg Gly Tyr Ile Thr Arg Tyr Phe Tyr Asn  
                                   105                                  110                                  115  
 Asn Gln Thr Lys Gln Cys Glu Arg Phe Lys Tyr Gly Gly Cys Leu Gly  
                                   120                                  125                                  130  
 15 Asn Met Asn Asn Phe Glu Thr Leu Glu Glu Cys Lys Asn Ile Cys Glu  
                                   135                                  140                                  145  
 Asp Gly Pro Asn Gly Phe Gln Val Asp Asn Tyr Gly Thr Gln Leu Asn  
 20          150                                  155                                  160  
 Ala Val Asn Asn Ser Leu Thr Pro Gln Ser Thr Lys Val Pro Ser Leu  
       165                                  170                                  175                                  180  
 25 Phe Glu Phe His Gly Pro Ser Trp Cys Leu Thr Pro Ala Asp Arg Gly  
                                   185                                  190                                  195  
 Leu Cys Arg Ala Asn Glu Asn Arg Phe Tyr Tyr Asn Ser Val Ile Gly  
                                   200                                  205                                  210  
 30 Lys Cys Arg Pro Phe Lys Tyr Ser Gly Cys Gly Gly Asn Glu Asn Asn  
                                   215                                  220                                  225  
 Phe Thr Ser Lys Gln Glu Cys Leu Arg Ala Cys Lys Lys Gly Phe Ile  
 35          230                                  235                                  240  
 Gln Arg Ile Ser Lys Gly Gly Leu Ile Lys Thr Lys Arg Lys Arg Lys  
       245                                  250                                  255                                  260  
 40 Lys Gln Arg Val Lys Ile Ala Tyr Glu Glu Ile Phe Val Lys Asn Met  
                                   265                                  270                                  275

&lt;210&gt; 3

&lt;211&gt; 912

45 &lt;212&gt; DNA

&lt;213&gt; Human

&lt;220&gt;

&lt;221&gt; CDS

50 &lt;222&gt; (1)..(912)

&lt;223&gt; TFPI mutant + signal peptide

&lt;220&gt;

&lt;221&gt; mat\_peptide

55 &lt;222&gt; (85)..(912)

&lt;223&gt; TFPI mutant mature peptide (single nucleotide polymorphism)

&lt;220&gt;

60 &lt;221&gt; mutation

&lt;222&gt; (536)

&lt;223&gt; mutation site: original C was replaced by T ( Pro by Leu)

<400> 3  
 atg att tac aca atg aag aaa gta cat gca ctt tgg gct tct gta tgc 48  
 Met Ile Tyr Thr Met Lys Lys Val His Ala Leu Trp Ala Ser Val Cys  
 5 -25 -20 -15

ctg ctg ctt aat ctt gcc cct gcc cct ctt aat gct gat tct gag gaa 96  
 Leu Leu Leu Asn Leu Ala Pro Ala Pro Leu Asn Ala Asp Ser Glu Glu  
 -10 -5 -1 1

10 gat gaa gaa cac aca att atc aca gat acg gat ttg cca cca ctg aaa 144  
 Asp Glu Glu His Thr Ile Ile Thr Asp Thr Asp Leu Pro Pro Leu Lys  
 5 10 15 20

15 ctt atg cat tca ttt tgt gca ttc aag gcg gat gat ggc cca tgt aaa 192  
 Leu Met His Ser Phe Cys Ala Phe Lys Ala Asp Asp Gly Pro Cys Lys  
 25 30 35

gca atc atg aaa aga ttt ttc ttc aat att ttc act cga cag tgc gaa 240  
 20 Ala Ile Met Lys Arg Phe Phe Phe Asn Ile Phe Thr Arg Gln Cys Glu  
 40 45 50

gaa ttt ata tat ggg gga tgt gaa gaa aat cag aat cga ttt gaa agt 288  
 25 Glu Phe Ile Tyr Gly Gly Cys Glu Glu Asn Gln Asn Arg Phe Glu Ser  
 55 60 65

ctg gaa gag tgc aaa aaa atg tgt aca aga gat aat gca aac agg att 336  
 Leu Glu Glu Cys Lys Lys Met Cys Thr Arg Asp Asn Ala Asn Arg Ile  
 70 75 80

30 ata aag aca aca ttg caa caa gaa aag cca gat ttc tgc ttt ttg gaa 384  
 Ile Lys Thr Thr Leu Gln Gln Glu Lys Pro Asp Phe Cys Phe Leu Glu  
 85 90 95 100

35 gaa gat cct gga ata tgt cga ggt tat att acc agg tat ttt tat aac 432  
 Glu Asp Pro Gly Ile Cys Arg Gly Tyr Ile Thr Arg Tyr Phe Tyr Asn  
 105 110 115

aat cag aca aaa cag tgt gaa cgt ttc aag tat ggt gga tgc ctg ggc 480  
 40 Asn Gln Thr Lys Gln Cys Glu Arg Phe Lys Tyr Gly Gly Cys Leu Gly  
 120 125 130

aat atg aac aat ttt gag aca ctg gaa gaa tgc aag aac att tgt gaa 528  
 45 Asn Met Asn Asn Phe Glu Thr Leu Glu Glu Cys Lys Asn Ile Cys Glu  
 135 140 145

gat ggt ctg aat ggt ttc cag gtg gat aat tat gga acc cag ctc aat 576  
 Asp Gly Leu Asn Gly Phe Gln Val Asp Asn Tyr Gly Thr Gln Leu Asn  
 150 155 160

50 gct gtg aat aac tcc ctg act ccg caa tca acc aag gtt ccc agc ctt 624  
 Ala Val Asn Asn Ser Leu Thr Pro Gln Ser Thr Lys Val Pro Ser Leu  
 165 170 175 180

55 ttt gaa ttt cac ggt ccc tca tgg tgt ctc act cca gca gac aga gga 672  
 Phe Glu Phe His Gly Pro Ser Trp Cys Leu Thr Pro Ala Asp Arg Gly  
 185 190 195

ttg tgt cgt gcc aat gag aac aga ttc tac tac aat tca gtc att ggg 720  
 60 Leu Cys Arg Ala Asn Glu Asn Arg Phe Tyr Tyr Asn Ser Val Ile Gly  
 200 205 210

aaa tgc cgc cca ttt aag tac agt gga tgt ggg gga aat gaa aac aat 768  
 Lys Cys Arg Pro Phe Lys Tyr Ser Gly Cys Gly Gly Asn Glu Asn Asn  
 215 220 225

5 ttt act tcc aaa caa gaa tgt ctg agg gca tgt aaa aaa ggt ttc atc 816  
 Phe Thr Ser Lys Gln Glu Cys Leu Arg Ala Cys Lys Lys Gly Phe Ile  
 230 235 240

10 caa aga ata tca aaa gga ggc cta att aaa acc aaa aga aaa aga aag 864  
 Gln Arg Ile Ser Lys Gly Gly Leu Ile Lys Thr Lys Arg Lys Arg Lys  
 245 250 255 260

15 aag cag aga gtg aaa ata gca tat gaa gaa att ttt gtt aaa aat atg 912  
 Lys Gln Arg Val Lys Ile Ala Tyr Glu Glu Ile Phe Val Lys Asn Met  
 265 270 275

<210> 4  
 20 <211> 304  
 <212> PRT  
 <213> Human

<400> 4  
 25 Met Ile Tyr Thr Met Lys Lys Val His Ala Leu Trp Ala Ser Val Cys  
 -25 -20 -15

Leu Leu Leu Asn Leu Ala Pro Ala Pro Leu Asn Ala Asp Ser Glu Glu  
 -10 -5 -1 1

30 Asp Glu Glu His Thr Ile Ile Thr Asp Thr Asp Leu Pro Pro Leu Lys  
 5 10 15 20

Leu Met His Ser Phe Cys Ala Phe Lys Ala Asp Asp Gly Pro Cys Lys  
 35 25 30 35

Ala Ile Met Lys Arg Phe Phe Phe Asn Ile Phe Thr Arg Gln Cys Glu  
 40 40 45 50

40 Glu Phe Ile Tyr Gly Gly Cys Glu Glu Asn Gln Asn Arg Phe Glu Ser  
 55 60 65

Leu Glu Glu Cys Lys Lys Met Cys Thr Arg Asp Asn Ala Asn Arg Ile  
 70 75 80

45 Ile Lys Thr Thr Leu Gln Gln Glu Lys Pro Asp Phe Cys Phe Leu Glu  
 85 90 95 100

Glu Asp Pro Gly Ile Cys Arg Gly Tyr Ile Thr Arg Tyr Phe Tyr Asn  
 50 105 110 115

Asn Gln Thr Lys Gln Cys Glu Arg Phe Lys Tyr Gly Gly Cys Leu Gly  
 120 125 130

55 Asn Met Asn Asn Phe Glu Thr Leu Glu Glu Cys Lys Asn Ile Cys Glu  
 135 140 145

Asp Gly Leu Asn Gly Phe Gln Val Asp Asn Tyr Gly Thr Gln Leu Asn  
 150 155 160

60 Ala Val Asn Asn Ser Leu Thr Pro Gln Ser Thr Lys Val Pro Ser Leu  
 165 170 175 180

Phe Glu Phe His Gly Pro Ser Trp Cys Leu Thr Pro Ala Asp Arg Gly  
 185 190 195  
 5 Leu Cys Arg Ala Asn Glu Asn Arg Phe Tyr Tyr Asn Ser Val Ile Gly  
 200 205 210  
 Lys Cys Arg Pro Phe Lys Tyr Ser Gly Cys Gly Gly Asn Glu Asn Asn  
 215 220 225  
 10 Phe Thr Ser Lys Gln Glu Cys Leu Arg Ala Cys Lys Lys Gly Phe Ile  
 230 235 240  
 Gln Arg Ile Ser Lys Gly Gly Leu Ile Lys Thr Lys Arg Lys Arg Lys  
 245 250 255 260  
 15 Lys Gln Arg Val Lys Ile Ala Tyr Glu Glu Ile Phe Val Lys Asn Met  
 265 270 275

20 <210> 5  
 <211> 5  
 <212> DNA  
 <213> Human  
 <220>  
 <221> misc\_feature  
 25 <222> (1)..(5)  
 <223> restriction site  
 <400> 5  
 actgg

5

30 <210> 6  
 <211> 20  
 <212> DNA  
 <213> Human  
 35 <221> primer\_bind  
 <222> (1)..(20)  
 <223> Primer  
 <400> 6  
 tctatttttaa ttggctgtat

20

40  
 <210> 7  
 <211> 5  
 <212> DNA  
 45 <213> Human  
 <221> misc\_feature  
 <222> (1)..(5)  
 <223> restriction site  
 <400> 7  
 50 cagtg

5

<210> 8  
 <211> 20  
 55 <212> DNA  
 <213> Human  
 <221> primer\_bind  
 <222> (1)..(20)  
 <223> Primer  
 60 <400> 8  
 gcatgataat agtttcctgg

20

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/06054

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K14/81 C12N15/52 C12N15/10 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 93 14122 A (NOVONORDISK AS) 22 July 1993 (1993-07-22) abstract page 4, line 12 -page 5, line 30 page 7, line 9-20 ---	1,10
A	WO 96 04378 A (CHIRON CORP) 15 February 1996 (1996-02-15) abstract page 4, line 1-12 page 5, line 10 -page 7, line 15 page 7, line 31 -page 9, line 21 --- -/--	1,4,10

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

10 December 1999

Date of mailing of the international search report

20/12/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Mateo Rosell, A.M.



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/06054

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	GIRARD T J ET AL., : "Functional significance of the Kunitz-type inhibitory domains of lipoprotein-associated coagulation inhibitor" NATURE, vol. 338, 1989, pages 518-520, XP000857349 cited in the application the whole document ---	1
A	SANDSET P M: "Tissue factor pathway inhibitor (TFPI): An update." HAEMOSTASIS, vol. 26 (SUPPL. 4), 1996, page 154-165 XP000862822 the whole document ---	1
P,X	KLEESIEK K ET AL., : " A first mutation in the human tissue factor pathway inhibitor gene encoding 'P151L!TFPI" BLOOD, vol. 92 (10), 15 November 1998 (1998-11-15), page 3976-7 XP000856946 the whole document ---	1-10
P,X	KLEESIEK K ET AL., : "The 536CfwdarwT transition in the human tissue factor pathway inhibitor (TFPI) gene is statistically associated with a higher risk for venous thrombosis." THROMBOSIS AND HAEMOSTASIS , vol. 82 (1), July 1999 (1999-07), page 1-5 XP000856948 the whole document ---	1-10
P,A	MIYATA T ET AL., : "C-399T polymorphism in the promoter region of human tissue factor pathway inhibitor (TFPI) gene does not change the plasma TFPI antigen level and does not cause venous thrombosis" THROMBOSIS AND HAEMOSTASIS , vol. 80 (2), August 1998 (1998-08), page 345-346 XP000856905 the whole document --- -/--	1,4-6,10

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 99/06054

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,A	<p>MOATTI D ET AL., : "Polymorphisms of the tissue factor pathway inhibitor (TFPI) gene in patients with acute coronary syndromes and in healthy subjects: Impact of the V264M substitution on plasma levels of TFPI."</p> <p>ARTERIOSCLEROSIS THROMBOSIS AND VASCULAR BIOLOGY, vol. 19 (4), April 1999 (1999-04), page 862-869 XP000856947 the whole document</p>	1,4-6,10

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 99/06054

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9314122 A	22-07-1993	AU 675926 B	27-02-1997
		AU 3346093 A	03-08-1993
		CA 2127246 A	22-07-1993
		CZ 9401644 A	15-12-1994
		EP 0621872 A	02-11-1994
		FI 943234 A	06-07-1994
		HU 70293 A	28-09-1995
		JP 7504891 T	01-06-1995
		NO 942549 A	07-09-1994
		NZ 246570 A	25-09-1996
		ZA 9300096 A	10-08-1993
WO 9604378 A	15-02-1996	US 5589359 A	31-12-1996
		AU 710535 B	23-09-1999
		AU 3150095 A	04-03-1996
		CA 2196290 A	15-02-1996
		EP 0776366 A	04-06-1997
		JP 10503375 T	31-03-1998
		US 5696088 A	09-12-1997
		US 5563123 A	08-10-1996